

4 Dimensional Phenotyping

Gareth Baynam^{1,2}, Mark Walters³, Peter Claes⁴, Stephanie Kung², Peter LeSouef², Hugh Dawkins^{5,6,7}, Matthew Bellgard⁸, Marta Girdea⁹, Peter Robinson¹⁰, Andreas Zankl¹¹, David Gillett³, Jack Goldblatt^{1,2}

¹Genetic Services of Western Australia, Princess Margaret & King Edward Memorial Hospitals, Perth, Western Australia, Australia, ²School of Paediatrics and Child Health, University of Western Australia, Perth, Western Australia, Australia ³Cranio-Maxillo-Facial Unit, Princess Margaret Hospital for Children, Perth, Western Australia, Australia, ⁴P. Claes, K.U. Leuven, Medical Imaging Research Centre, Faculty of Engineering, Leuven, Belgium, ⁵Office of Population Health Genomics, Department of Health, Perth, Western Australia, Australia, ⁶Centre for Population Health Research, Curtin Health Innovation Research Institute, Curtin University of Technology, Perth, Western Australia, Australia, ⁷School of Pathology and Laboratory Medicine, University of Western Australia, Perth, Western Australia, Australia, ⁸Center for Comparative Genomics, Murdoch University; Department of Computer Science, University of Toronto, Ontario, Canada, ⁹Institut für Medizinische Genetik und Humangenetik, Charité – Universitätsmedizin, Berlin, Germany, ¹⁰The University of Queensland, UQ Centre for Clinical Research, Queensland, Australia

Analysis of the clinical phenotype: history, examination and investigation, is the physician's daily work¹. Documentation of patient phenotypes has become particularly to clarify genotype–phenotype relationships that are increasingly being investigated by the use of high throughput molecular technologies. To improve the clinical utility of emerging -omics technologies, such as next generation sequencing, there is a parallel and increasing need for efficient, and preferably objective and standardised, documentation of the clinical phenotype. This will require time-efficient tools for ascertaining phenotypic information, incorporating this information into clinical work flows, and for integrating with -omics analyses. This presentation will focus on aspects of phenotyping that are of particular relevance to rare diseases and which have implications for common diseases. Firstly, for illustrative purposes, it will discuss 3-dimensional facial analysis: a fine scale, objective, scalable, portable, cost efficient, non-invasive and non-irradiating phenotypic assessment². Secondly, it will describe the “4th dimension” of bed-side capture of standardized phenotypic information, workflow integration and emerging decision support tools.

Robinson PN. 2012. Deep phenotyping for precision medicine. *Hum Mutat* 33(5):777–780

Baynam G, Walters M, Claes P, Kung S, LeSouef P, Dawkins H, Gillett D, Goldblatt J. 2013. The facial evolution: looking backward and moving forward. *Hum Mutat* 34(1):14–22